



Melting curve anomalies of ^3He and ^4He

Both isotopes show an **unusual minimum** in their melting curves for **very different reasons**

^4He : very shallow (hardly to see) minimum at 0.8 K because the **phonon entropy** is higher in the solid phase

^3He : pronounced minimum at 0.32 K because the **nuclear spin entropy** is higher in the solid phase



Clausius–Clapeyron equation

$$\left. \frac{\partial p}{\partial T} \right|_{\text{meltingcurve}} = \frac{S_l - S_s}{V_l - V_s}$$

If $V_l > V_s$ and $S_s > S_l$ the **slope** of the **melting curve** becomes **negative**



Here the example of ^3He



Liquid ^3He is a Landau liquid
more in Chapter 3




a) Specific heat

^4He : first measurements 1926 by Kamerlingh Onnes and Dana
(rise at T_λ neglected)
later Keesom and Clausius **discovery of phase transition** at T_λ at **2.17 K**


Explanation not before 1938:

first idea: new **crystalline** phase
Model of liquid crystal
but X-ray scattering results





^3He : **Discovery of phase transition** with NMR by Richarson, Lee, Osheroff before specific heat measurements
(also wrong interpretation: phase transition in solid ^3He)






b) Density

^4He : maximum at T_λ

^3He : smooth monotone temperature dependence






c) Latent heat

evaporation of helium

line: from vapor pressure measurement and Clausius-Claperyron equation

^4He : kink at T_λ

^3He : smooth temperature dependence





2.1 Experimental Observations


a) Boiling

at boiling point: liquid \longleftrightarrow dense classical gas


at lambda point $T_\lambda = 2.17 \text{ K}$ boiling **ceases abruptly!**
transition from **He-I** to **He-II**



$T > T_\lambda$



$T \sim T_\lambda$



$T < T_\lambda$



b) Viscosity

measurement: **flow** through **thin capillaries**

Hagen-Poiseuille law

$$\dot{V} = \frac{\pi r^4}{8} \frac{1}{\eta} \frac{\Delta p}{L}$$

volume rate




flow velocity

$$v = \dot{V} / (\pi r^2)$$


$$v \propto r^2 \Delta p$$

viscosity



Experimental results:

- ▶ v **independent** of pressure
- ▶ v **increasing** with **decreasing** diameter





Conclusion:

$$\eta_{\text{He-II}} < 10^{-3} \eta_{\text{He-I}} < 10^{-2} \eta_{\text{H}_2\text{O}}$$

→ discovery of superfluidity 1938


Kapitza
Allen, Misener

Question: $\eta_{\text{He-II}} = 0$?

→ persistent flow experiments 1965

Reppy, Mehl
Zimmermann

Torus with **fine** powder and He



slowly breaking, He-II rotates further

→ measurement of angular velocity


Results: **angular velocity constant over many hours**

$$\eta_{\text{He-II}} < 10^{-11} \eta_{\text{He-I}} \quad \eta_{\text{He-II}} \stackrel{!}{=} 0 \quad \text{within measurement error}$$




Measurements with **3 standard methods**


Capillaries, Slits



Rotary Viscosimeter



Oscillating Disc





➔ results are **seemingly** completely **inconsistent**



1 Pa/s = 10 P (Poise)



c) Beaker Experiments




- ▶ helium flows over the rim of beakers
- ▶ helium flows with constant rate independent of level difference
- ▶ flow can be reversed at equal rate



- ▶ detailed measurement with thin neck \rightarrow small ΔV \rightarrow large Δh
- ▶ oscillations are observed when level equalizes \rightarrow not damped (in special cases)
- \rightarrow persistent flow



d) Thermomechanical Effect



Very thin capillary (super leak)



course of the experiment

1. pressures and temperatures are equal
2. pressure increases in A
3. helium flows to B
4. temperature in A increases and drops in B

$$\rightarrow \Delta p \rightarrow T_A^{\uparrow}, T_B^{\downarrow} \rightarrow T_A > T_B$$

▶ Mass flow is connected with heat transport

▶ $\Delta p \curvearrowright \Delta T$, but heat flow is in opposite direction of mass flow



- ▶ **heating** of helium inside vessel ΔT Δp helium shoots out at the top
- ▶ **stationary heights** up to **30 cm**, have been observed!