


Neutrino scattering (Part 1)

Neutrino nucleon scattering played an important role to understand the weak interaction and also to resolve the proton and neutron structure in DIS:



Fixed target configuration:




$$\begin{aligned}
 s &= (E_\nu + m_N)^2 - E_\nu^2 \\
 &= 2m_N E_\nu - m_N^2 \\
 &\approx 2m_N E_\nu \quad \text{for } E_\nu \gg m_N
 \end{aligned}$$

Fundamental process: ν -Quark scattering, discussed in the following.


A few words on neutrino beams

$\bar{\nu}_\mu$ beams are produced from high-energy proton beams



Focusing is done with a so called "magnetic horn" (see slide)

v-Quark scattering



Matrix element $v/ q^2 \ll m_W^2 : \frac{1}{q^2 - m_W^2} \rightarrow \frac{1}{m_W^2}$

$$-iM_{fi} = \left[-i \frac{g}{\sqrt{2}} \bar{u}(p_3) \gamma^\mu \frac{1}{2} (1 - \gamma^5) u(p_1) \right] \cdot \frac{i g_{WV}}{m_W^2} \left[-i \frac{g}{\sqrt{2}} \bar{u}(p_4) \gamma^\nu \frac{1}{2} (1 - \gamma^5) u(p_2) \right]$$

(effective theory) $\left\{ \frac{1}{q^2 - m_W^2} \rightarrow \frac{1}{m_W^2} \right\}$

- $\frac{1}{2}(1 - \gamma^5)$ projects out the LH spinor components
- γ^μ coupling ensures that only $\bar{u}_L \gamma^\mu u_L$ term contribute.

If we neglect the particle masses: $u_L = u_{\downarrow}$ i.e. chiral states = helicity state

$$\Rightarrow M_{fi} = \frac{g^2}{2m_W^2} \cdot \left[\bar{u}_{\downarrow}(p_3) \delta_{\mu\nu} u_{\downarrow}(p_1) \right] \left[\bar{u}_{\downarrow}(p_4) \gamma^\mu u_{\downarrow}(p_2) \right]$$

Using the Dirac-Pauli representation of the γ -matrices and of the spinors (see exercise sheets 2 and 3):

$$u_{\downarrow}(p_1) = \sqrt{E} \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix} \quad (\varphi_1=0, \theta_1=0)$$

$$u_{\downarrow}(p_2) = \sqrt{E} \begin{pmatrix} -1 \\ 0 \\ 1 \\ 0 \end{pmatrix} \quad (\varphi_2=\pi, \theta_2=\pi)$$

$$u_{\downarrow}(p_3) = \sqrt{E} \begin{pmatrix} -s \\ 0 \\ s \\ -c \end{pmatrix} \quad (\varphi_3=0, \theta_3=\theta)$$

$$u_{\downarrow}(p_4) = \sqrt{E} \begin{pmatrix} -c \\ -s \\ c \\ s \end{pmatrix} \quad (\varphi_4=\pi, \theta_4=\pi-\theta)$$

in CMS, $m_i \approx 0$
 $|\vec{p}_i| = |\vec{E}_i| = E$
 $\hat{S} = 4E^2$
 with $s = \sin \theta$
 $c = \cos \theta$
 see slide

one finds: $j_e^\mu = \bar{u}_{\downarrow}(p_3) \gamma^\mu u_{\downarrow}(p_1) = 2E (c, s, -is, c)$

$j_q^\mu = \bar{u}_{\downarrow}(p_4) \gamma^\mu u_{\downarrow}(p_2) = 2E (c, -s, -is, -c)$

$$\Rightarrow M_{fi} = \frac{g^2}{2m_W^2} \cdot 4E^2 (c^2 + s^2 + s^2 + c^2) = \frac{g^2}{m_W^2} \cdot \hat{S}$$

(For the cross section we need to average M_{fi} over all initial spin configurations and sum of all possible final spin configs:
Spin averaging + summation:

Initial: ν is always LH; neutrinos can be LH, RH however only LH contribute. Averaging $\rightarrow \frac{1}{2}$

final: only LH configuration contribute.

$$\Rightarrow \langle |M_{fi}|^2 \rangle = \frac{1}{2} \cdot |M_{fi}|^2 = \frac{1}{2} \left(\frac{g^2}{m_W^2} \hat{s} \right)^2$$

With formula for diff. cross section:

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2} \frac{1}{s} \langle |M_{fi}|^2 \rangle \Rightarrow \frac{d\sigma_{\nu q}}{d\Omega_{\mu}} = \frac{1}{64\pi^2} \frac{1}{\hat{s}} \cdot \frac{1}{2} \left(\frac{g^2}{m_W^2} \right)^2 \hat{s}^2$$

$$\Rightarrow \frac{d\sigma}{d\Omega_{\mu}} = \frac{G_F^2}{4\pi^2} \cdot \hat{s} \quad (\text{in CMS system}).$$

\uparrow
 with $\frac{G_F}{\sqrt{2}} = \frac{g^2}{8m_W^2}$

Total cross section (integration over 4π):

$$\sigma_{\nu q} = \frac{G_F^2}{\pi} \cdot \hat{s} \quad \left. \begin{array}{l} \text{not really measurable.} \\ \text{What can be measured} \\ \text{is } \nu N \text{ cross section.} \end{array} \right\}$$

For very large \hat{s} , i.e. at large E_{ν} result violates unitarity:


Used "effective theory" is not valid anymore. We

should have used "full theory" with propagator $\frac{1}{q^2 - m_W^2}$

\rightarrow additional factor $\frac{1}{s^2}$ which cures the unitarity problem.

\rightarrow Fermi's theory w/ 4-fermion interaction is not valid anymore and we need a theory with "propagating W's" (i.e. SM)

$\bar{\nu}_\mu$ -quark scattering



$$M_{fi} = \frac{g^2}{2m_W^2} \cdot \left[\bar{\nu}_\mu(p_3) \gamma_\mu \nu_\mu(p_1) \right] \left[\bar{u}_d(p_4) \gamma^\mu u_d(p_2) \right]$$

$j_{\ell, \mu}$ $j_{q, \mu}$

using the anti-spinors (see slide):

$$\nu_\uparrow(p_1) = \sqrt{E} \begin{pmatrix} 0 \\ -1 \\ 0 \\ 1 \end{pmatrix}$$

$\varphi_1 = 0 \quad \theta_1 = 0$

$$\nu_\uparrow(p_3) = \sqrt{E} \begin{pmatrix} s \\ -c \\ s \\ c \end{pmatrix}$$

$\varphi_3 = \pi \quad \theta_3 = \theta$

$$s = \sin \frac{\theta}{2}$$

$$c = \cos \frac{\theta}{2}$$

one finds in analysis: $M_{fi} = \frac{g^2}{m_W^2} \cdot \frac{1}{2} (1 + \cos \theta) \hat{s}$

or $\langle |M_{fi}|^2 \rangle = \frac{1}{2} \left(\frac{g^2}{m_W^2} \right)^2 \cdot \frac{1}{4} (1 + \cos \theta)^2 \hat{s}^2$

diff. cross section:

$$\frac{d\sigma_{\bar{\nu}q}}{d\Omega_{\nu}} = \frac{G_F^2}{4\pi^2} \cdot \hat{s} \cdot \frac{1}{4} (1 + \cos \theta)^2 \quad (\text{in CMS})$$


integration $\int_{\Omega} \dots = \frac{4\pi}{3}$

$$\Rightarrow \sigma_{\bar{\nu}q} = \frac{G_F^2}{4\pi^2} \hat{s} \cdot \frac{4\pi}{3} = \frac{G_F^2 \cdot \hat{s}}{3\pi} = \frac{1}{3} \sigma_{\nu q}$$

At the first view, this result is surprising.
 (Becomes clear if we analyze the spin configurations)


spin configurations of incoming massless particles:

νq - scattering





total spin = 0
flat $\cos\theta$ - distribution
of outgoing particles

$\bar{\nu} q$ - scattering



total spin = 1
angular distribution falls
 $\sim (1 + \cos\theta)^2$



$\bar{\nu}$ - antiquark scattering


In case of νq and $\bar{\nu} q$ one can easily convince oneself that the roles are exchanged: anti RH antiquarks couple to W.

$$\Rightarrow \begin{cases} \frac{d\sigma_{\nu q}}{d\Omega_{\mu}} = \frac{G_F^2}{4\pi^2} \cdot \hat{s} \cdot \frac{1}{4} (1 + \cos\theta)^2 \\ \frac{d\sigma_{\bar{\nu} q}}{d\Omega_{\mu}} = \frac{G_F^2}{4\pi^2} \cdot \hat{s} \end{cases}$$

Measured $\nu N / \bar{\nu} N$ cross section ratio (s. slide) shows ~ 0.5 instead of $1/3 \Rightarrow$ evidence that nucleons also contain anti-quarks!

Neutral current " v-scattering

One of the predictions of the SM is the NC (i.e. Z exchange) in v-scattering process:





Z couples to LH and RH quarks with different strengths:

$$C_L^q = \frac{1}{2} (C_V^q + C_A^q)$$

$$C_R^q = \frac{1}{2} (C_V^q - C_A^q)$$

Reminder:	$C_V = T_3 - 2Q \sin^2 \theta_W$	$C_A = T_3$	$\sin^2 \theta_W \approx 0.231$
u-type	$T_3 = +1/2$	$Q = +2/3$	$C_V = \frac{1}{2} - \frac{4}{3} \sin^2 \theta_W \approx +0.19$
d-type Quarks	$T_3 = -1/2$	$Q = -1/3$	$C_V = -\frac{1}{2} + \frac{2}{3} \sin^2 \theta_W \approx -0.35$
			$C_A = +1/2$
			$C_L \approx 0.35$
			$C_R \approx -0.15$
			$\approx -0.42 \approx +$

We should thus discuss to different spin configurations:



Instead of redoing the calculation one can "adapt" the CC-result.

$$\frac{d\sigma_{\nu q}^{NC}}{d\Omega_{\nu q}} = \frac{G_F^2}{4\pi^2} \cdot \hat{s} \cdot \left[(g_L^q)^2 + \frac{1}{4} (g_R^q)^2 (1 + \cos \theta)^2 \right]$$

$\Rightarrow \sigma_{\nu q}^{NC} = \frac{G_F^2}{\pi} \hat{s} \left[(g_L^q)^2 + \frac{1}{3} (g_R^q)^2 \right]$ *How does it change? for $\bar{\nu}q$ scattering?*

Remarks: - Observation of v-scattering w/o N^\pm confirms Z exchange predicted by SM

- $R = \frac{\sigma_{\nu q}^{NC}}{\sigma_{\nu q}^{CC}} \sim (g_L^q)^2 + \frac{1}{3} (g_R^q)^2 \sim f(\sin^2 \theta_W)$
 \rightarrow see slide: Discovery of NC